



ENTERIC VIRUSES IN WATER AS A RISK FACTOR IN A MEGACITY

Ana C Espinosa
Universidad Nacional Autónoma de México,
Mexico City
acespino@ecologia.unam.mx

Background

94% of all diarrhea cases attributable to the environment

88% associated with risk factors

→ contaminated drinking
water/poor sanitation (WHO,
2006)





Coliform bacteria → most common indicator fecal water contamination

Bacterial indicators do not correlate number and viability of enteric viruses

Viruses > resistant than bacteria environmental factors and water treatment

Infective doses of viruses < than bacteria


Developing countries ->
epidemic information
incidence of viral
gastroenteritis unknown,
incomplete or not updated



Table 1. Infective doses of different pathogen groups

Pathogen	Infective dose		
Enteric viruses	1 PFU		
Protozoa	25-100 cysts/oocystes		
Fecal bacteria	>100 CFU		

Alonso et al, 2006.



Rotavirus



Giardia intestinalis (cyst)



Escherichia coli

Mexico City Metropolitan Area (MEGACITY)

> 22 million inhabitants

High water demand drinking and others uses

Wastewater production (68 m³/s)

Defficient/inadequate wastewater treatment (<10%)




Concern about drinking water quality

Inadequate reuse of treated wastewater for irrigation purposes

Specific case study

→ Universidad Nacional
Autónoma de México (UNAM)





Multidisciplinary program → managment, use, reuse of water, conduced at UNAM

Main goals

- ✓ Safe drinking water for university community, representative area South of Mexico City
- ✓ Promote safe reuse of treated wastewater



Cycle 2009: cold-dry, warm-dry and rainy season

Presence and quantity of indicator bacteria, enterovirus, adenovirus, coliphage in groundwater, wastewater, and treated wastewater reused for recreational areas irrigation





Groundwater

Reuse treated wastewater



UNAM wastewater treatment plant







Sample concentration by ultrafiltration (polysulphone hollow fibers)



Membrane filtration for fecal coliform (FC) and fecal enterococcus (FE) quantification in m-FC and KF media



qPCR for enterovirus (EV) and adenovirus (AdV) genome quantification


- ✓EV conserved region for enterovirus 5'NTR
- ✓ AdV conserved region for AdV 40 and 41, L4 region






Double agar layer for coliphage F-RNA specific quantification using *Escherichia coli* K12 host bacteria






Results

Wastewater showed highest counts for all indicators: FC, FE, EV, AdV and coliphage








Fig 1. Detection of indicator bacteria (A), enteric viruses and coliphage (B) in wastewater for three periods

FC and FE not detected in treated wastewater→ adequate bacteria remotion. Enteric viruses and coliphage detected




Fig 2. Enteric virus (EV, AdV) and coliphage in treated wastewater for two periods (wastewater not treated in rainy season).




Fig 3. Indicator bacteria (A), enteric viruses and coliphage (B), in treated wastewater reused for irrigation during two periods (no irrigation during rainy season)

Table 2. Indicator bacteria (CFU/100 mL), enteric viruses (genomes/mL), coliphage (PFU/mL) in groundwater for three periods (2009)

Period -	Indicator				
	FC	FE	AdV	EV	Coliphages
cold-dry	1	1	31	2	<1
warm-dry	<1	<1	<1	<1	23
rainy	<1	1	<1	<1	20

Conclusions

Although wastewater treatment achieved to remove FC and FE, reused water for irrigation showed bacterial re-growth or re-contamination

FC, FE counts and enteric virus in reused water for irrigation hazards to the University community exposed in green areas

Based on microbial indicators -> groundwater from University campus+disinfection adequate source for drinking water

Monitoring enteric viruses in urban wastewater and reused water → necessary and useful as a risk assessment tool for the exposed population

Water managment must consider risk assessment studies in order to take better decisions to avoid or decrease health impacts

Acknowledgments

Managment, use and reuse of water program (*PUMAGUA*) UNAM-IINGEN

CONACyT project No. 60577

Participants

Dra. Marisa Mazari-Hiriart

Dr. Fernando González Villarreal

Biol. Miguel Atl Silva Magaña

Biol. Alejandra Fonseca Salazar

Biol. Rosa Solano Ortíz

Gamsahamnida

